

New Jersey College Immunization Reminder for Healthcare Providers 2024-2025

Check patient immunization records to make sure they are ready for Back to Campus! *†‡§

CD	CDC Recommended Schedule	Doses
Hepatitis B	≥ 0 months	3 doses (2 doses may be acceptable based on age indication and vaccine)
Measles, Mumps, Rubella (MMR)	≥ 12 months	2 doses at least 28 days apart
Meningococcal (ACWY) ‡ <i>New</i> as of June 15, 2020	16 -18 years	1 dose
Meningococcal (B) ‡ <i>New</i> as of June 15, 2020	16-23 years at increased risk; those not at high risk may receive vaccine based on shared clinical decision making.	2 or 3 doses depending on vaccine; boosters as recommended

* NJ has minimum immunization requirements for attending institutions of Higher Education which may differ from the recommendations listed above. For optimal protection, NJDOH recommends following the CDC Recommended Immunization Schedules. Refer to N.J.A.C. 8:57-6 for New Jersey specific immunization requirements, including medical and religious exemptions.

† Additional vaccines may be recommended based on age, occupation, lifestyle, or risk factors. Click [Adult Immunization Schedule – Healthcare Providers | CDC](#) for the CDC Recommended and Catch-Up Adult Immunization Schedules.

‡ For more information on the Higher Education Meningococcal Requirement, click [here](#).

§ A list of currently accepted COVID vaccines can be found on the CDC [website](#). For specific information, visit [Clinical Guidance for COVID-19 Vaccination | CDC](#).

New Jersey College Immunization Reminder for Healthcare Providers 2024-2025

Check patient immunization records to make sure they are ready for Back to Campus! *†‡§

	CDC Recommended Schedule	Doses
COVID-19 §	≥ 6 months	Check most recent recommendation
Hepatitis A	≥ 12 months	2 or 3 doses depending on vaccine
Human papillomavirus (HPV)	9-26 years Persons 27 through 45 may receive vaccine based on shared clinical decision making.	2 or 3 doses depending on age at initial vaccination and risk factors
Influenza	≥ 6 months	1 dose annually
Mpox†	Persons with an additional risk factor or indication	2 doses at least 28 days apart
Pneumococcal Vaccine	≥ 65 years 19-64 years with additional risk factor or indication	Varies depending on indication and product
Respiratory Syncytial Virus (RSV)†	RSV vaccine for pregnant person or monoclonal antibody for babies younger than 8 months during the baby's first RSV season	1 dose
Shingles	≥ 50 years ≥ 19 with additional risk factor or indication	2 doses
Tetanus-Diphtheria-Pertussis (Tdap)	11- 12 years; Children who receive Tdap ≥ 10 years do not need the routine Tdap at age 11 – 12 years	1 dose; Td or Tdap booster every 10 years
Varicella (Chicken Pox)	≥ 12 months	2 doses

* NJ has minimum immunization requirements for attending institutions of Higher Education which may differ from the recommendations listed above. For optimal protection, NJDOH recommends following the CDC Recommended Immunization Schedules. Refer to N.J.A.C. 8:57-6 for New Jersey specific immunization requirements, including medical and religious exemptions.

† Additional vaccines may be recommended based on age, occupation, lifestyle, or risk factors. Click [Adult Immunization Schedule – Healthcare Providers | CDC](#) for the CDC Recommended and Catch-Up Adult Immunization Schedules.

‡ For more information on the Higher Education Meningococcal Requirement, click [here](#).

§ A list of currently accepted COVID vaccines can be found on the CDC [website](#). For specific information, visit [Clinical Guidance for COVID-19 Vaccination | CDC](#).